

Subject: The Lower Anogenital Tract Terminology Standardization Project for HPV-Associated Lesions (LAST) Guideline Update: Articles in Evidence Tables
(Literature search dates: 1/1/2012 – 10/23/2024)

1. Alshenawy HA. Evaluation of p16, human papillomavirus capsid protein L1 and Ki-67 in cervical intraepithelial lesions: Potential utility in diagnosis and prognosis. *Pathology, Research & Practice*. 2014;210(12):916-921. PMID: 25149503 (RefID 2049)
2. Aslani FS, Safaei A, Pourjabali M, Momtahan M. Evaluation of Ki67, p16 and CK17 markers in differentiating cervical intraepithelial neoplasia and benign lesions. *Iranian Journal of Medical Sciences*. 2013;38(1):15-21. PMID: 23645953 (RefID 3306)
3. Cavalcante DM, Linhares IM, Pompeu MM, Giraldo PC, Eleuterio J. The utility of p16 INK4a and Ki-67 to identify high-grade squamous intraepithelial lesion in adolescents and young women. *Indian J Pathol Microbiol*. 2012;55(3):339-342. PMID: 23032827 (RefID 223)
4. Clark JL, Lu D, Kalir T, Liu Y. Overdiagnosis of HSIL on cervical biopsy: Errors in p16 immunohistochemistry implementation. *Hum Pathol*. 2016;55:51-56. PMID: 27134110 (RefID 1003)
5. Clinton LK, Miyazaki K, Ayabe A, et al. The LAST guidelines in clinical practice: Implementing recommendations for p16 use. *Am J Clin Pathol*. 2015;144(6):844-849. PMID: 26572990 (RefID 61)
6. da Costa LBE, Triglia RM, Andrade L. P16INK4a, cytokeratin 7, and Ki-67 as potential markers for low-grade cervical intraepithelial neoplasia progression. *J Low Genit Tract Dis*. 2017;21(3):171-176. PMID: 28403025 (RefID 950)
7. da Costa LB, Triglia RdM, Franca Junior MC, Andrade LA. P16(INK) (4a) expression as a potential marker of low-grade cervical intraepithelial neoplasia progression. *APMIS*. 2015;123(3):185-189. PMID: 25471335 (RefID 973)
8. Dcunha NJ, Khandekar M, Puranik S. Role of p16 immunohistochemistry in cervical intraepithelial neoplasia. *J Clin Diagn Res*. 2020;14(3):1-5. <https://doi.org/10.7860/JCDR/2020/41707.13549> (RefID 3161)
9. De Lima TM, De Azevedo Focchi GR, De Almeida BC, Napoli Belfort-Mattos P, Heinke T, De Gois Speck NM. Expression of CK7 and CDKN2 in cervical intraepithelial neoplasia and correlation with clinical outcome. *Anticancer Res*. 2018;38(12):6673-6681. PMID: 30504376 (RefID 2001)
10. de Melo FL, Lancellotti CL, da Silva MA. Expression of the immunohistochemical markers p16 and Ki-67 and their usefulness in the diagnosis of cervical intraepithelial neoplasms. *Rev Bras Ginecol Obstet*. 2016;38(2):82-87. PMID: 26883858 (RefID 1978)
11. de Sam Lazaro S, Newbill CP, Berlin M, Morgan TK. P16 staining of cervical biopsies may decrease the frequency of unnecessary loop electrosurgical excision procedures. *J Low Genit Tract Dis*. 2016;20(3):201-206. PMID: 26855146 (RefID 978)

12. Ding L, Song L, Zhao W, et al. Predictive value of p16INK4a, Ki-67 and ProExC immuno-qualitative features in LSIL progression into HSIL. *Exp Ther Med*. 2020;19(4):2457-2466. PMID: 32256722 (RefID 787)
13. Ezaldein H, Lott JP, McNiff JM, et al. Grading of atypia in genital skin lesions: Routine microscopic evaluation and use of p16 immunostaining. *J Cutan Pathol*. 2015;42(8):519-526. PMID: 25951050 (RefID 1841)
14. Ferraz DLF, Dos Santos ES, Pedroso CM, et al. Histopathological risk assessment in multisite epithelial dysplasia: A meta-analysis. *Oral Dis*. 2024. PMID: 38946217 (RefID 5445)
15. Hissong E, Cimic A, Leeman A, Pirog EC. Utility of Stathmin-1 as a novel marker in evaluating anal intraepithelial neoplasia (AIN). *Appl Immunohistochem Mol Morphol*. 2019;27(2):134-139. PMID: 28777150 (RefID 3415)
16. Howitt BE, Nucci MR, Drapkin R, Crum CP, Hirsch MS. Stathmin-1 expression as a complement to p16 helps identify high-grade cervical intraepithelial neoplasia with increased specificity. *Am J Surg Pathol*. 2013;37(1):89-97. PMID: 23211296 (RefID 405)
17. Kremer WW, Vink FJ, van Zummeren M, et al. Characterization of cervical biopsies of women with HIV and HPV co-infection using p16INK4a, Ki-67 and HPV E4 immunohistochemistry and DNA methylation. *Mod Pathol*. 2020;33(10):1968-1978. PMID: 32249820 (RefID 2704)
18. Krishnamurti U, Mohammad M, Monsrud A, et al. Diagnosing anal squamous intraepithelial lesions with and without p16: An interobserver variability study. *J Low Genit Tract Dis*. 2020;24(1):69-74. PMID: 31860579 (RefID 2290)
19. Lee S, Kim H, Kim H, Kim C, Kim I. The utility of p16INK4a and Ki-67 as a conjunctive tool in uterine cervical lesions. *Korean J Pathol*. 2012;46(3):253-260. PMID: 23110011 (RefID 221)
20. Lee YY, Song SY, Do IG, et al. Dynamin 2 expression as a biomarker in grading of cervical intraepithelial neoplasia. *Eur J Obstet Gynecol Reprod Biol*. 2012;164(2):180-184. PMID: 22959143 (RefID 2170)
21. Liao GD, Sellors JW, Sun HK, et al. P16INK4a immunohistochemical staining and predictive value for progression of cervical intraepithelial neoplasia grade 1: A prospective study in China. *Int J Cancer*. 2014;134(7):1715-1724. PMID: 24105727 (RefID 954)
22. Liu DQ, Li FF, Zhang JB, et al. Increased RIPK4 expression is associated with progression and poor prognosis in cervical squamous cell carcinoma patients. *Sci Rep*. 2015;5:11955. PMID: 26148476 (RefID 1335)
23. Liu Y, McCluggage WG, Darragh TM, et al. P16 immunoreactivity correlates with morphologic diagnosis of HPV-associated anal intraepithelial neoplasia: A study of 1000 biopsies. *Am J Surg Pathol*. 2021;45(11):1573-1578. PMID: 34231547 (RefID 988)
24. Liu Y, Blakely M, Sigel K, et al. Biomarker p16 predicts progression risk of anal low-grade squamous intraepithelial lesions. *AIDS*. 2018;32(16):2309-2316. PMID: 30005024 (RefID 2831)

25. Liu Y, McCluggage WG, Darragh TM, et al. Classifying anal intraepithelial neoplasia 2 based on LAST recommendations. *Am J Clin Pathol*. 2021;155(6):845-852. PMID: 33210115 (RefID 2678)
26. Lukic A, Rossi S, Frega A, et al. Prognostic role of immunohistochemical overexpression of the p16 protein in women under the age of 35 and diagnosed with HSIL (CIN2) subjected to "cervix sparing" excision. *Eur Rev Med Pharmacol Sci*. 2021;25(3):1261-1273. PMID: 33629296 (RefID 748)
27. Mandal R, Ghosh I, Banerjee D, et al. Correlation between p16/Ki-67 expression and the grade of cervical intraepithelial neoplasias. *Int J Gynecol Pathol*. 2020;39(4):384-390. PMID: 32515923 (RefID 2428)
28. Meserve E, Berlin M, Mori T, Krum R, Morgan TK. Reducing misclassification bias in cervical dysplasia risk factor analysis with p16-based diagnoses. *J Low Gen Tract Dis*. 2014;18(3):266-272. PMID: 24633165 (RefID 652)
29. Mills AM, Dirks DC, Poulter MD, Mills SE, Stoler MH. HR-HPV E6/E7 mRNA in situ hybridization. Validation against PCR, DNA in situ hybridization, and p16 immunohistochemistry in 102 samples of cervical, vulvar, anal, and head and neck neoplasia. *Am J Surg Pathol*. 2017;41(5):607-615. PMID: 28403015 (RefID 3673)
30. Mills AM, Paquette C, Terzic T, Castle PE, Stoler MH. CK7 immunohistochemistry as a predictor of CIN1 progression: A retrospective study of patients from the quadrivalent HPV vaccine trials. *Am J Surg Pathol*. 2017;41(2):143-152. PMID: 27680604 (RefID 2682)
31. Miralpeix E, Genoves J, Maria Sole-Sedeno J, et al. Usefulness of p16INK4a staining for managing histological high-grade squamous intraepithelial cervical lesions. *Mod Pathol*. 2017;30(2):304-310. PMID: 27739439 (RefID 236)
32. Montero S, Sandigo-Saballos I, Tom C, Lee H. Poor acceptance of the revised classification of premalignant anal lesions following the last standardized project. *Am Surg*. 2024;90(10):2419-2423. PMID: 38641876 (RefID 4621)
33. Nooij LS, Dreef EJ, Smit VT, van Poelgeest MI, Bosse T. Stathmin is a highly sensitive and specific biomarker for vulvar high-grade squamous intraepithelial lesions. *J Clin Pathol*. 2016;69(12):1070-1075. PMID: 27226646 (RefID 404)
34. Pacchiarotti A, Galeotti S, Bellardini P, et al. Impact of p16(INK4a) immunohistochemistry staining on interobserver agreement on the diagnosis of cervical intraepithelial neoplasia. *Am J Clin Pathol*. 2014;141(3):367-373. PMID: 24515764 (RefID 1368)
35. Roberts JM, Jin F, Thurlow JK, et al. High reproducibility of histological diagnosis of human papillomavirus-related intraepithelial lesions of the anal canal. *Pathology*. 2015;47(4):308-313. PMID: 25938361 (RefID 84)
36. Roma AA, Liu X, Patil DT, Xie H, Allende D. Proposed terminology for anal squamous lesions: Its application and interobserver agreement among pathologists in academic and community hospitals. *Am J Clin Pathol*. 2017;148(1):81-90. PMID: 28605400 (RefID 718)
37. Salim RC, Lima TM, Heinke T, et al. Cervical squamous intraepithelial lesions grades 1 and 2: Value of an alternative morphological classification and of p16 immunohistochemistry in the prediction of clinical

38. Sripathi S, Mamatha M, Akarapu JD, Pyla RD. Histomorphological evaluation of cervical punch biopsies with the aid of p16 INK4a and Ki 67. Eur J Mol Clin Med. 2022;9(6):1269-1286. (RefID 5316)
39. Stoler MH, Wright TC, Jr., Ferenczy A, et al. Routine use of adjunctive p16 immunohistochemistry improves diagnostic agreement of cervical biopsy interpretation: Results from the CERTAIN study. Am J Surg Pathol. 2018;42(8):1001-1009. PMID: 29697437 (RefID 510)
40. Tao AS, Zuna R, Darragh TM, et al. Interobserver reproducibility of cervical histology interpretation with and without p16 immunohistochemistry. Am J Clin Pathol. 2024;162(2):202-209. PMID: 38527169 (RefID 4915)
41. Thrall MJ. Effect of lower anogenital squamous terminology recommendations on the use of p16 immunohistochemistry and the proportion of high-grade diagnoses in cervical biopsy specimens. Am J Clin Pathol. 2016;145(4):524-530. PMID: 27124943 (RefID 2148)
42. Umphress B, Sanchez B, Paintal A, Nayar R, Maniar KP. Utility of CK7 versus p16 as a prognostic biomarker in CIN 2. Am J Surg Pathol. 2018;42(4):479-484. PMID: 29438168 (RefID 231)
43. van Baars R, Griffin H, Wu Z, et al. Investigating diagnostic problems of CIN1 and CIN2 associated with high-risk HPV by combining the novel molecular biomarker panHPV4 with p16INK4a. Am J Surg Pathol. 2015;39(11):1518-1528. PMID: 26379150 (RefID 1289)
44. Wright TC, Jr., Stoler MH, Ferenczy A, et al. The certain study results: Adjunctive p16 immunohistochemistry use in cervical biopsies according to LAST criteria. Am J Surg Pathol. 2021;45(10):1348-1356. PMID: 34232603 (RefID 2785)
45. Xing Y, Wang C, Wu J. Expression of geminin, p16, and Ki67 in cervical intraepithelial neoplasm and normal tissues. Medicine (Baltimore). 2017;96(26):e7302. PMID: 28658133 (RefID 1996)
46. Yang J, Elliott A, Hoffa AL, Herring N, Houser PM. Potential influence of p16 immunohistochemical staining on the diagnosis of squamous cell lesions in cervical biopsy specimens: Observation from cytologic-histologic correlation. Cancer Cytopathol. 2018;126(12):1003-1010. PMID: 30307707 (RefID 803)

Disclaimer

The information, data, and draft recommendations provided by the College of American Pathologists are presented for informational and public feedback purposes only. The draft recommendations and supporting documents will be removed on September 10, 2025. The draft recommendations along with the public comments received and completed evidence review will be reassessed by the expert panel in order to formulate the final recommendations. These draft materials should not be stored, adapted, or redistributed in any manner.