

Template for Reporting Results of *HER2 (ERBB2)* Biomarker Testing of Specimens From Patients With Adenocarcinoma of the Stomach or Esophagogastric Junction

Template web posting date: June 2014

Authors

Angela N. Bartley, MD, FCAP

Department of Pathology, St. Joseph Mercy Hospital, Ann Arbor, MI

Jessi Christ, CTR

Sanford Health Medical Center, Fargo, ND

Patrick Fitzgibbons, MD, FCAP

Department of Pathology, St. Jude Medical Center, Fullerton, CA

Stanley R. Hamilton, MD, FCAP

Division of Pathology and Laboratory Medicine, University of Texas MD Anderson Cancer Center, Houston, TX

Sanjay Kakar, MD, FCAP

Department of Pathology, University of California San Francisco and the Veterans Affairs Medical Center, San Francisco, CA

Manish A. Shah, MD

Weill Cornell Medical College, New York, NY

Laura H. Tang, MD, PhD, FCAP

Department of Pathology, Memorial Sloan-Kettering Cancer Center, New York, NY

Megan L. Troxell MD, PhD

Oregon Health and Science University, Portland, OR

For the Members of the Cancer Biomarker Reporting Committee, College of American Pathologists

© 2014 College of American Pathologists (CAP). All rights reserved.

The College does not permit reproduction of any substantial portion of these templates without its written authorization. The College hereby authorizes use of these templates by physicians and other health care providers in reporting results of biomarker testing on patient specimens, in teaching, and in carrying out medical research for nonprofit purposes. This authorization does not extend to reproduction or other use of any substantial portion of these templates for commercial purposes without the written consent of the College.

The CAP also authorizes physicians and other health care practitioners to make modified versions of the templates solely for their individual use in reporting results of biomarker testing for individual patients, teaching, and carrying out medical research for non-profit purposes.

The CAP further authorizes the following uses by physicians and other health care practitioners, in reporting on surgical specimens for individual patients, in teaching, and in carrying out medical research for non-profit purposes: (1) **Dictation** from the original or modified templates for the purposes of creating a text-based patient record on paper, or in a word processing document; (2) **Copying** from the original or modified templates into a text-based patient record on paper, or in a word processing document; (3) The use of a **computerized system** for items (1) and (2), provided that the template data is stored intact as a single text-based document, and is not stored as multiple discrete data fields.

Other than uses (1), (2), and (3) above, the CAP does not authorize any use of the templates in electronic medical records systems, pathology informatics systems, cancer registry computer systems, computerized databases, mappings between coding works, or any computerized system without a written license from the CAP.

Any public dissemination of the original or modified templates is prohibited without a written license from the CAP.

The College of American Pathologists offers these templates to assist pathologists in providing clinically useful and relevant information when reporting results of biomarker testing. The College regards the reporting elements in the templates as important elements of the biomarker test report, but the manner in which these elements are reported is at the discretion of each specific pathologist, taking into account clinician preferences, institutional policies, and individual practice.

The College developed these templates as educational tools to assist pathologists in the useful reporting of relevant information. It did not issue them for use in litigation, reimbursement, or other contexts. Nevertheless, the College recognizes that the templates might be used by hospitals, attorneys, payers, and others. The College cautions that use of the templates other than for their intended educational purpose may involve additional considerations that are beyond the scope of this document.

The inclusion of a product name or service in a CAP publication should not be construed as an endorsement of such product or service, nor is failure to include the name of a product or service to be construed as disapproval.

CAP Gastric HER2 Biomarkers Template Revision History

Version Code

The definition of the version code can be found at www.cap.org/cancerprotocols.

Version: GastricHER2Biomarkers 1.0.0.0

Summary of Changes

This is a new template.

Gastric HER2 Biomarker Reporting Template

Template web posting date: June 2014

Completion of the template is the responsibility of the laboratory performing the biomarker testing and/or providing the interpretation. When both testing and interpretation are performed elsewhere (eg, a reference laboratory), synoptic reporting of the results by the laboratory submitting the tissue for testing is also encouraged to ensure that all information is included in the patient's medical record and thus readily available to the treating clinical team.

STOMACH/ESOPHAGOGASTRIC JUNCTION

Select a single response unless otherwise indicated.

Note: Use of this template is optional.

+ RESULTS

+ HER2 (by immunohistochemistry)

- + ___ Negative (score 0)
- + ___ Negative (score 1+)
- + ___ Equivocal (score 2+)
- + ___ Positive (score 3+)
- + ___ Indeterminate (explain): _____

+ HER2 (ERBB2) (by in situ hybridization)

- + ___ Negative (not amplified)
- + ___ Positive (amplified)
- + ___ Indeterminate (explain): _____

+ Number of observers: _____

+ Number of invasive cancer cells counted: _____

- + ___ Using dual-probe assay
 - + Average number of HER2 (ERBB2) signals per cancer cell: _____
 - + Average number of CEP17 signals per cancer cell: _____
 - + HER2 (ERBB2):CEP17 ratio: _____

- + ___ Using single-probe assay
 - + Average number of HER2 (ERBB2) signals per cancer cell: _____

+ METHODS

+ HER2 (by immunohistochemistry)

- + ___ US Food and Drug Administration (FDA) cleared (specify test/vendor): _____
- + ___ Laboratory-developed test

+ Data elements preceded by this symbol are not required.

+ Primary Antibody

- + ___ 4B5
- + ___ HercepTest™
- + ___ A0485
- + ___ SP3
- + ___ CB11
- + ___ Other (specify): _____

+ **HER2 (ERBB2) (by in situ hybridization)**

- + ___ FDA cleared (specify test/vendor): _____
- + ___ Laboratory-developed test (specify probe): _____

Gene names should follow recommendations of The Human Genome Organisation (HUGO) Nomenclature Committee (www.genenames.org; accessed February 10, 2015).

All reported gene sequence variations should be identified following the recommendations of the Human Genome Variation Society (www.hgvs.org/mutnomen/; accessed February 10, 2015).

Explanatory Notes

HER2 (ERBB2) is a proto-oncogene located on chromosome 17 that encodes a 185-kd tyrosine kinase receptor belonging to the epidermal growth factor receptor (EGFR) family whose phosphorylation initiates signaling pathways that lead to cell division, proliferation, differentiation, and apoptosis.¹⁻³ The Human Genome Organisation (HUGO) Nomenclature Committee (HGNC) has designated *ERBB2* as the approved symbol and CD340, HER-2, HER2, and NEU as synonyms (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=3430). *HER2* gene product is expressed in normal epithelial cells, and amplification and/or overexpression of this gene has been reported in up to 30% of breast cancers⁴ and in 9% to 27% of patients with gastric cancer. Overexpression in stomach cancer varies with histologic type (intestinal type greater than diffuse type) and differentiation (moderately differentiated greater than poorly differentiated).⁵

For patients with inoperable locally advanced, recurrent, or metastatic adenocarcinoma of the stomach or esophagogastric junction for whom trastuzumab (Herceptin) is under consideration for therapy, assessment for tumor *HER2* overexpression using immunohistochemistry (IHC) or in situ hybridization (ISH) is recommended by the National Comprehensive Cancer Network (NCCN).⁵ Results of an open-label, international, phase 3 randomized controlled trial in 2010 (Trastuzumab for Gastric Cancer [ToGA]) showed that the anti-*HER2* humanized monoclonal antibody trastuzumab is effective in prolonging survival compared with chemotherapy alone in patients with *HER2*-positive adenocarcinoma of the stomach and the esophagogastric junction.⁶ *HER2 (ERBB2)* appears to be an important prognostic factor in gastric cancer, although the literature is conflicting, and not all studies have shown an association between *HER2* overexpression and poor prognosis.^{4,7} Clinical trials with antibodies to *HER2* in gastric cancer patients are in progress.

HER2 (ERBB2) status is assessed by testing either biopsy or surgical resection specimens. IHC evaluates membranous protein expression of cancer cells. Both intensity and percentage of immunoreactive cancer cells is assessed with scores ranging from 0 to 3+ (Table 1). ISH, which encompasses fluorescence in situ hybridization (FISH), chromogenic in situ hybridization (CISH), and silver-enhanced in situ hybridization (SISH), identifies the presence or absence of gene amplification. Some assays use a single *HER2 (ERBB2)* probe to determine the number of *HER2 (ERBB2)* gene copies present, but most assays include a chromosome enumeration probe (CEP17) to determine the ratio of *HER2 (ERBB2)* signals to copies of chromosome 17. ISH has been used to verify IHC-equivocal cases.⁵ *HER2*-positive gastric cancer has been defined as IHC 3+ or ISH positive in the USA and Japan, and IHC 3+ or 2+ with ISH positivity in Europe.^{4,8} In the US, the FDA has approved trastuzumab in association with chemotherapy for metastatic gastric cancer utilizing the eligibility criteria of the ToGA trial, limited to patients with a score of IHC 3+ or 2+ and ISH positivity. No significant survival benefit was seen for patients who were IHC 0 or 1+ and FISH positive.⁹

HER2 protein expression is more heterogeneous in gastric cancers than in breast cancers.^{7,8,10} The completeness of membrane staining required for positivity in breast cancers is infrequent in gastric adenocarcinomas, which often exhibit a basolateral staining pattern. Detection of *HER2 (ERBB2)* gene amplification by FISH is similar to that in breast cancer according to the American Society of Clinical Oncology/College of American Pathologists (ASCO/CAP) 2013 guidelines, in which *HER2 (ERBB2)* amplification is defined as *HER2 (ERBB2)*:CEP17 ratio of ≥ 2 .¹¹ Hoffman et al developed a four-tier scoring system for IHC (Table 1), also used in the ToGA trial, for gastric cancer by using the assessment area cutoff of at least 10% stained tumor cells for resection specimens and a small cluster of cells (≥ 5 neoplastic cells) for biopsy specimens.⁷ The NCCN guidelines recommend that assessment for *HER2* status should be performed first using immunohistochemistry following the modified scoring system used in the ToGA trial. A score of 0 or 1+ is considered to be negative for *HER2* expression. A score of 2+ is considered equivocal and should be confirmed with FISH or other in situ hybridization techniques. The NCCN panel recommends FISH only for cases with IHC 2+, although some institutions routinely perform

both IHC and FISH on all cases. The guidelines recommend trastuzumab with chemotherapy only for patients with IHC 3+ and IHC 2+ with evidence of *HER2 (ERBB2)* amplification by ISH (*HER2 (ERBB2):CEP17* ratio ≥ 2). Trastuzumab is not recommended if the IHC score is 0 or 1+.⁵

Table 1. Criteria Used in the ToGA Trial⁶ for Scoring HER2 Expression by Immunohistochemistry (IHC) in Gastric and Esophagogastric Junction Adenocarcinoma

HER2 IHC Score	HER2 IHC Pattern in Surgical Specimen	HER2 IHC Pattern in Biopsy Specimen	HER2 Expression Assessment
0	No reactivity or membranous reactivity in <10% of cancer cells	No reactivity or no membranous reactivity in any cancer cell	Negative by IHC
1+	Faint or barely perceptible membranous reactivity in $\geq 10\%$ of cancer cells; cells are reactive only in part of their membrane	Cancer cell cluster* with a faint or barely perceptible membranous reactivity irrespective of percentage of cancer cells positive	Negative by IHC
2+	Weak to moderate complete, basolateral or lateral membranous reactivity in >10% of tumor cells	Cancer cell cluster* with a weak to moderate complete, basolateral, or lateral membranous reactivity irrespective of percentage of cancer cells positive	Equivocal by IHC
3+	Strong complete, basolateral or lateral membranous reactivity in $\geq 10\%$ of cancer cells	Cancer cell cluster* with a strong complete basolateral, or lateral membranous reactivity irrespective of percentage of cancer cells positive	Positive

* Cancer cell cluster consisting of ≥ 5 neoplastic cells

References

1. Akiyama T, Sudo C, Ogawara H, Toyoshima K, Yamamoto T. The product of the human c-erbB-2 gene: a 185-kilodalton glycoprotein with tyrosine kinase activity. *Science*. 1986;232:1644-1646
2. Ménard S, Pupa SM, Campiglio M, Tagliabue E. Biologic and therapeutic role of HER2 in cancer. *Oncogene*. 2003;22:6570-6578.
3. Lui L, Wu N, Li J. Novel targeted agents for gastric cancer. *J Hematol Oncol*. 2012;5:31.
4. Boku N. HER2 positive gastric cancer. *Gastric Cancer*. 2014;17(1):1-12.
5. Ajani JA, Bentrem DJ, Besh S, et al. Gastric cancer, version 2.2013: featured updates to the NCCN Guidelines. *J Natl Compr Canc Netw*. 2013;11:531-546.
6. Bang YJ, Van Cutsem E, Feyereislova A, et al. Trastuzumab in combination with chemotherapy versus chemotherapy alone for treatment of HER2-positive advanced gastric or gastro-oesophageal junction cancer (ToGA): a phase 3, open-label, randomized controlled trial. *Lancet*. 2010;376:687-697.
7. Hofmann M, Stoss O, Shi D, et al. Assessment of a HER2 scoring system for gastric cancer: results from a validation study. *Histopathology*. 2008;52:797-805.
8. Albarello L, Pecciarini L, Doglioni C. HER2 testing in gastric cancer. *Adv Anat Pathol*. 2011;18:53-9.
9. Bang YJ. Advances in the management of HER2-positive advanced gastric and gastroesophageal junction cancer. *J Clin Gastroenterol*. 2012;46:637-648.
10. Pirrelli M, Caruso ML, Di Maggio M, Armentano R, Valentini AM. Are biopsy specimens predictive of HER2 status in gastric cancer patients? *Digest Dis Sci*. 2013;58:397-404.

11. Wolff AC, Hammond MEH, Hicks DG, et al. Recommendations for human epidermal growth factor receptor 2 testing in breast cancer: American Society of Clinical Oncology/College of American Pathologists clinical practice guideline update. *Arch Pathol Lab Med*. 2014;138(2):241-256.